


HERBICIDE RECROPPING RESTRICTIONS

The 2022 growing season had varied precipitation accumulation across the province, not only due to heavy spring flooding but throughout the season. In late October, the Available Water Holding Capacity was measured from 0 – 120 cm depth in over 100 locations across Manitoba to develop the map below. While most of the province was experiencing at or slightly below water holding capacity, there are areas that were significantly drier. With ample moisture over the 2022 growing season, there should be little concern about herbicide carryover. However, there can still be issue, regardless of soil moisture, so become familiar with recropping restrictions in each commodity.

Recropping restrictions are implemented to prevent crop damage following applications of residual herbicides. The following table is a guide for corn producers to identify any fields that should not be planted to the crop in a specific time frame, following the application of the correlating herbicide.

Continues on next page »



Canada

Manitoba

Figure 1: Manitoba Fall Soil Moisture Survey via Manitoba Agriculture: Percent of Available Water Holding Capacity in the

0 – 120 cm Depth. https://www.gov.mb.ca/agriculture/environment/soil-management/manitoba-fall-soil-moisture-survey.html



Figure 2: Reflex (fomesafen) herbicide carryover damage in corn plant.



HERBICIDE RECROPPING RESTRICTIONS

NOTE: Herbicide residue breakdown occurs through several processes in and on the soil. Chemical reactions, soil microbes and leaching are just a few methods of breakdown and are strongly favoured by warm, moist soil conditions (not achieved during spring thaw). If the applied field did not receive adequate precipitation during the growing season, the herbicide residual will remain, requiring recropping restrictions to be lengthened. Contact the associated company for more information and recommendations.

Product	Company	Recropping Restrictions for Corn
Altitude FX3	BASF Canada	2 years**
Clopyralid/MCPA (+/- fluroxypyr)		1 year**
Dicamba/Fluroxypyr		2 years
Distinct	BASF Canada	30 days
Fierce	Valent Canada/Nufarm Agriculture	7 days
Flexstar GT	Syngenta Canada	10 months** (RRV only)
Fluroxypyr		1 year*
lmazamox/lmazethapyr		1 year*
Imazethapyr		2 years
Momentum	Loveland Products Canada	2 years
Muster	FMC Corporation	2 years+
Nicosulfuron		10 months
Olympus	Bayer	NR+
Permit WG	Gowan Canada	30 days
Quinclorac		NR+
Signal FSU	Nufarm Agriculture	2 years
Smoulder	BASF Canada	11 months
Thifen:triben (2:1) + fluroxypyr		2 years
Thifen:Triben (25:25) + fluroxypyr		12 months
Travallas	FMC Corporation	10 months
Trifluralin		1 year**

 $\ensuremath{\mathbf{NR:}}$ May be planted or replanted the year of application. No recropping restrictions. NR+: No re-cropping restrictions indicated for field corn; farmer should conduct a field bioassay to ensure crop safety and contact the company for more information.

DISCLAIMER: The recropping intervals listed may not be sufficient to prevent crop injury during periods of below average precipitation. The information provided is only a guide, Manitoba Crop Alliance is not liable for any decisions made based on this information. Always refer to the product label for application details and precautions.

REFERENCES: Guide to Field Crop Protection, Manitoba Agriculture and Resource Development. 2023



mbcropalliance.ca hello@mbcropalliance.ca P: 204.745.6661







^{*:} May not be supported for all products; refer to product label for details.

^{**:} May be longer depending on use rates, region, soil type, environment, time of application

^{+:} Farmer should conduct a bio-assay to ensure crop safety and contact the company for more information